



International Journal of Innovative Technologies in Social Science

e-ISSN: 2544-9435

Operating Publisher
SciFormat Publishing Inc.
ISNI: 0000 0005 1449 8214

2734 17 Avenue SW,
Calgary, Alberta, T3E0A7,
Canada
+15878858911
editorial-office@sciformat.ca

ARTICLE TITLE	THE IMPACT OF ARTIFICIAL INTELLIGENCE ON HEARING HEALTHCARE: DIAGNOSTICS AND THERAPEUTIC INNOVATIONS
----------------------	---

DOI	https://doi.org/10.31435/ijitss.4(52).2026.6531
------------	---

RECEIVED	17 August 2026
-----------------	----------------

ACCEPTED	01 October 2026
-----------------	-----------------

PUBLISHED	07 October 2026
------------------	-----------------

LICENSE	
----------------	--



The article is licensed under a **Creative Commons Attribution 4.0 International License**.

© The author(s) 2026.

This article is published as open access under the Creative Commons Attribution 4.0 International License (CC BY 4.0), allowing the author to retain copyright. The CC BY 4.0 License permits the content to be copied, adapted, displayed, distributed, republished, or reused for any purpose, including adaptation and commercial use, as long as proper attribution is provided.

THE IMPACT OF ARTIFICIAL INTELLIGENCE ON HEARING HEALTHCARE: DIAGNOSTICS AND THERAPEUTIC INNOVATIONS

Wiktoria Balińska (Corresponding Author, Email: wiktoria.balinska@stud.umed.lodz.pl)
Medical University of Lodz, Poland
ORCID ID: 0009-0002-2615-7236

Aleksandra Marek
Medical University of Lodz, Poland
ORCID ID: 0009-0002-3391-5082

Jagoda Rogowska
Medical University of Lodz, Poland
ORCID ID: 0009-0004-5995-5802

Jakub Wąsik
Medical University of Lodz, Poland
ORCID ID: 0009-0002-7145-8118

ABSTRACT

Background: Artificial Intelligence (AI) is progressively transforming otolaryngology, enhancing diagnostics and treatment precision. AI-powered tools improve hearing aid personalization, assist in cochlear implant mapping, and analyze audiometric and imaging data for accurate diagnosis. Machine learning also advances tinnitus management through personalized sound therapy. This review aims to explore the current applications of AI in audiology and otolaryngology, highlighting its benefits, limitations, and future directions in clinical practice.

Methods: A comprehensive literature search was performed in November 2024 across PubMed, Embase, Medline, and Scopus using the terms: Artificial Intelligence in Hearing Loss, AI in Diagnostics of Hearing Loss, AI in Cochlear Implants, and Personalized Treatment for Hearing Loss using AI. Duplicates and unrelated publications were excluded, and full-text articles were assessed for eligibility with disagreements resolved by consensus.

Results: The final selection included clinical trials, observational studies, and case reports, focusing on AI's diagnostic capabilities and hearing intervention advancements. These studies were categorized into three main themes: "AI in Audiological Diagnostics," "Advancing Hearing Loss Treatment with AI," and "Challenges and Ethical Considerations."

Conclusions: Artificial intelligence has the potential to revolutionize audiology by improving diagnostics, optimizing hearing devices, and enhancing patient outcomes. Ongoing collaboration among audiologists, data scientists, and engineers, incorporating ethical considerations, will be key to fully realizing AI's impact on hearing and balance disorder treatment.

KEYWORDS

Artificial Intelligence (AI), Otolaryngology, Audiology, Machine Learning, Hearing Diagnostics, Hearing Loss

CITATION

Balińska, W., Marek, A., Rogowska, J., & Wąsik, J. (2026). The impact of artificial intelligence on hearing healthcare: Diagnostics and therapeutic innovations. *International Journal of Innovative Technologies in Social Science*, 4(52). [https://doi.org/10.31435/ijitss.4\(52\).2026.6531](https://doi.org/10.31435/ijitss.4(52).2026.6531)

COPYRIGHT

© The author(s) 2026. This article is published as open access under the **Creative Commons Attribution 4.0 International License (CC BY 4.0)**, allowing the author to retain copyright. The CC BY 4.0 License permits the content to be copied, adapted, displayed, distributed, republished, or reused for any purpose, including adaptation and commercial use, as long as proper attribution is provided.

Introduction

Artificial Intelligence (AI), broadly defined as the development of computer systems designed to replicate complex human intellectual processes, has evolved significantly since its theoretical inception in the 1950s. The field is currently experiencing its third and most impactful boom, driven by exponential increases in computational power and the availability of vast amounts of digital data. In the broader scope of medicine, and particularly within the specialized field of otolaryngology, AI is demonstrating increasingly impressive diagnostic capabilities. For instance, in the realm of otoscopy, AI algorithms—predominantly based on deep learning and convolutional neural networks—have been deployed to detect conditions such as otitis media directly from tympanic membrane images. In many instances, these automated diagnostic tools perform at a level that equals or even surpasses the diagnostic accuracy of human clinicians. Furthermore, AI substantially enhances the evaluation of complex radiological imaging, including high-resolution computed tomography (CT) and magnetic resonance imaging (MRI) of the temporal bone. By automating image segmentation and highlighting microscopic anatomical structures, AI improves the diagnostic precision for intricate pathological conditions such as otosclerosis and vestibular schwannoma.

Beyond diagnostics, AI is making profound contributions to therapeutic interventions and surgical planning. Predictive machine learning models are now utilized to forecast clinical outcomes for patients experiencing sudden sensorineural hearing loss, as well as to predict the functional success rates of tympanoplasty surgeries based on preoperative variables. Additionally, AI optimizes the daily performance of auditory prostheses; modern hearing aids utilize machine learning to dynamically adjust amplification parameters in noisy environments, vastly improving speech recognition. AI also assists clinicians with the complex task of cochlear implant mapping, helping to predict post-implantation outcomes even in anatomically complex scenarios, such as severe cochlear malformations. Alongside these specialized tools, emerging trends feature generative AI and large language models, such as ChatGPT. These platforms are increasingly being explored for their potential to provide preliminary medical advice and assist in patient education, though their clinical reliability, accuracy, and tendency to generate hallucinated information require rigorous further validation before integration into standard practice [1].

Within the specific domains of audiology and phoniatrics, artificial intelligence is being deployed in a multitude of innovative ways. A prominent example is the transformative advancement in modern hearing aid technology. AI-powered hearing devices are now capable of continuously analyzing complex environmental acoustic scenes in real-time. By rapidly identifying whether a user is in a quiet room, a reverberant hall, or a noisy restaurant, the device automatically adjusts its digital signal processing settings to optimize the user's auditory experience. This high degree of personalization significantly improves speech intelligibility in highly challenging, signal-to-noise ratio-poor environments, directly addressing one of the most persistent and frustrating challenges for individuals suffering from hearing impairment [2].

In the domain of audiological diagnostics, sophisticated AI models have been developed to rapidly process and analyze complex audiometric data. These systems can distinguish between different phenotypes and etiologies of hearing loss with exceptionally high degrees of accuracy, minimizing human error in audiogram interpretation. Such models prove to be particularly valuable in resource-limited or remote clinical settings, where timely access to highly trained audiologists and specialized diagnostic equipment may be severely restricted. Leveraging this technology, AI-powered mobile audiometry platforms and smartphone-based applications have been successfully deployed to conduct large-scale population screening programs in a manner that is both time-efficient and highly cost-effective, thus democratizing access to early hearing healthcare [3].

The application of AI extends robustly into the advanced imaging techniques routinely utilized in audiological assessments, such as high-resolution CT and MRI scans of the brain and auditory pathways. These AI-enhanced analytical tools assist radiologists and otolaryngologists in identifying subtle structural abnormalities, ranging from congenital cochlear malformations to the early detection of vestibular schwannomas (acoustic neuromas). For example, a recent study demonstrated how advanced convolutional neural networks (CNNs) could be trained to autonomously analyze complex imaging data, identifying optimal cochlear implant candidates and assessing surgical feasibility with a level of accuracy that closely mirrors expert human assessment [4,5].

In the challenging area of tinnitus management—a condition notorious for its subjective nature and diverse etiology—AI has recently emerged as a highly valuable tool for developing individualized therapeutic strategies. Advanced machine learning algorithms possess the capability to analyze vast arrays of patient data, successfully identifying hidden patterns in tinnitus perception and predicting the potential clinical effectiveness of various established treatments, such as targeted sound therapy or cognitive behavioral therapy (CBT). Furthermore, novel AI-driven applications can precisely analyze the specific frequency and masking profile of an individual patient's tinnitus, subsequently generating highly tailored, dynamic sound therapy tracks designed to therapeutically target and modulate specific hyperactive auditory pathways in the brain [6].

Despite its vast transformative potential, the practical integration of AI into routine clinical otolaryngology presents several significant challenges. Foremost among these is the absolute necessity for large, diverse, and meticulously annotated clinical datasets required to train robust, generalizable machine learning algorithms that perform consistently across different patient demographics. Additionally, the "black box" nature of deep learning necessitates improved interpretability of AI models, ensuring that clinicians can transparently understand the reasoning behind an AI-generated diagnosis or recommendation [7]. Ethical considerations are similarly paramount, particularly those concerning the strict maintenance of patient data privacy, cybersecurity, and the nuances of informed consent in automated healthcare. Effectively addressing these multifaceted challenges requires deep, ongoing collaboration among practicing clinicians, academic researchers, legal experts, and healthcare policymakers to establish standardized, robust guidelines for the ethical and safe use of AI in otolaryngological practice [8].

Ultimately, artificial intelligence holds extraordinary, paradigm-shifting potential to revolutionize overall healthcare delivery; however, its tangible, day-to-day impact on routine medical practice has thus far remained somewhat limited. The primary reasons for this slow adoption stem from widespread misconceptions among both medical professionals and patients regarding exactly how these algorithms function, often fueled by an overestimation of AI's current autonomous capabilities. There are profound, systemic concerns regarding the perceived lack of direct human oversight, the fear of complete automation replacing nuanced clinical judgment, and the potentially severe medical and legal liabilities associated with algorithmic errors, misdiagnoses, or inherent dataset biases. Overcoming these barriers will require rigorous education, transparent clinical trials, and a fundamental paradigm shift—moving away from viewing AI as an autonomous replacement, and instead fully embracing it as an assistive, "human-in-the-loop" tool that augments the indispensable elements of human empathy and final clinical authority [9].

The aim of this study is to explore the transformative role of artificial intelligence in otolaryngology, particularly in improving diagnostic accuracy and optimizing interventions for hearing loss. By analyzing current advances and challenges, this review aims to highlight the potential of artificial intelligence in clinical practice and emphasize the importance of interdisciplinary collaboration in shaping future approaches to the management of audiological disorders.

Materials and Methods

Search Strategy

A systematic literature search was conducted in November 2024 across four electronic databases: PubMed, Embase, Medline (via Ovid), and Scopus. The following key terms and their combinations were used, adapted to each database's syntax with Boolean operators (AND/OR) and Medical Subject Headings (MeSH) where applicable:

- Population: Hearing loss, auditory disorders, cochlear implants
- Intervention: Artificial intelligence, machine learning, deep learning, neural networks
- Outcomes: Diagnostic accuracy, treatment efficacy, clinical applications

Sample Search String (PubMed):

Copy ("Artificial Intelligence"[Mesh] OR "Machine Learning"[Mesh] OR "Deep Learning"[Mesh] OR "Neural Networks"[Mesh]) AND ("Hearing Loss"[Mesh] OR "Deafness"[Mesh] OR "Cochlear Implants"[Mesh] OR "Auditory Diseases"[Mesh]) AND ("Diagnosis"[Mesh] OR "Therapeutics"[Mesh] OR "Treatment Outcome"[Mesh])

No date or language restrictions were applied to avoid excluding seminal studies. Gray literature (e.g., conference abstracts, preprints) was searched via Google Scholar and ProQuest to mitigate publication bias.

Eligibility Criteria

- Inclusion:
 - Original research (randomized controlled trials, cohort studies, case-control studies) or meta-analyses.
 - Studies evaluating AI applications in hearing loss diagnosis (e.g., audiometry, imaging) or treatment (e.g., hearing aids, cochlear implants).
 - Human subjects; no age restrictions.
- Exclusion:
 - Animal studies, editorials, or reviews without primary data.
 - Studies lacking explicit AI methodology (e.g., non-machine learning statistical analyses).
 - Non-English articles without available translations.

Study Selection Process

1. **Deduplication:** Records were imported into EndNote X20, and duplicates were removed using automated and manual checks.

2. **Title/Abstract Screening:** Two independent reviewers (W.B. and A.M.) screened titles/abstracts against eligibility criteria using Rayyan QCRI. Conflicts were resolved by a third reviewer (J.R.).

3. **Full-Text Review:** Potentially eligible studies underwent full-text review by the same reviewers. Discrepancies (<5% of cases) were resolved via consensus.

4. **PRISMA Flowchart:** The selection process was documented in a PRISMA flowchart (Figure 1.), detailing excluded studies and reasons (e.g., "irrelevant population").




Fig. 1. Identification of studies via databases and other sources

Data Extraction and Quality Assessment

The data extraction process was conducted using a standardized template designed to capture three key categories of information. First, study characteristics were recorded, including author names, publication year, country of origin, study design, and sample size. Second, detailed information about AI methodologies was extracted, encompassing the type of model employed (such as convolutional neural networks [CNN] or support vector machines [SVM]), the composition and source of training datasets, and validation metrics (including area under the curve [AUC], sensitivity, specificity, and accuracy values). Third, clinical outcome measures were documented, focusing on diagnostic performance metrics, treatment success rates, and comparative analyses between AI systems and clinician performance.

For quality assessment, three validated tools were utilized according to study design: the QUADAS-2 instrument was applied to evaluate risk of bias in diagnostic accuracy studies, the ROBINS-I tool was used for non-randomized intervention studies, and the Cochrane Risk of Bias Tool was employed for randomized controlled trials. This comprehensive approach ensured systematic evaluation of methodological quality across all included studies while maintaining consistency with best practices in evidence synthesis. Two independent reviewers performed both data extraction and quality assessment, with any discrepancies resolved through discussion and consensus. The standardized template facilitated direct comparison of findings across studies while preserving study-specific details essential for contextual interpretation of results.

Statistical Analysis

Where applicable, meta-analysis was planned using RevMan 5.4 (random-effects model, I^2 for heterogeneity). For narrative synthesis, findings were categorized by AI application (diagnostics/treatment) and outcome.

Ethical Considerations

No ethical approval was required as only publicly available data were analyzed. Patient data from included studies were anonymized and aggregated.

Results

A systematic literature search identified a total of 864 studies from which duplicates were excluded. After reviewing the abstracts, 340 full-text articles were assessed for eligibility based on predefined inclusion and exclusion criteria. This process included a detailed assessment of the relevance of each study to the topic of artificial intelligence in hearing loss diagnosis and intervention. After a comprehensive full-text review and resolution of any disagreements between reviewers, 48 articles were found to meet the inclusion criteria. Of these 48 studies, 21 were considered to provide a sufficient level of evidence based on the quality of their methodology and relevance to the research question.

The selected studies included a mixture of clinical trials, observational studies and case reports, reflecting a broad spectrum of research on the applications of artificial intelligence in hearing loss. These studies contributed valuable insights into the diagnostic capabilities of AI, its potential in aural rehabilitation and current advances in AI-assisted listening interventions. The results of these studies are summarised below, highlighting key trends, challenges and outcomes for AI integration in otolaryngology. It was decided that the results would be collated and presented in two categories: 'AI in Audiological Diagnostics' and 'Advancing Hearing Loss Treatment with AI'. This scheme provides a clear and understandable division of the manuscript for the reader, reflecting the management of the entire therapeutic process of the patient.

AI in Audiological Diagnostics

Hearing Screening & Risk Prediction

AI applications in hearing loss detection demonstrate significant diagnostic capabilities across multiple domains. Automated analysis of audiometric data enables accurate classification of hearing loss types (conductive, sensorineural, mixed), with particular utility in resource-limited settings [10]. Population screening programs utilizing AI algorithms show effectiveness in identifying at-risk individuals through analysis of demographic and clinical parameters [10]. Predictive models incorporating patient history and audiometric data provide actionable risk assessments for clinical decision-making [11].

In tympanic membrane image analysis, AI systems achieve 80-90% accuracy in detecting pathologies such as perforations and effusions, exceeding physician performance (75%) [12]. A Danish algorithm demonstrates 92.6% accuracy for otitis media detection, though lacking subtype differentiation [13]. Subsequent development by Zafer et al. yielded a model capable of distinguishing between acute otitis media, otitis media with effusion, and chronic suppurative otitis media [14]. These technical capabilities represent measurable advances in diagnostic accuracy for otologic conditions.

Temporal Bone CT Imaging

AI demonstrates significant improvements in temporal bone CT interpretation, overcoming limitations of traditional manual methods. Current automated segmentation algorithms achieve comparable accuracy to manual approaches while substantially reducing processing time [15]. The W-Net architecture developed by Lv et al. provides particularly efficient analysis of complex temporal bone anatomy [5].

For specific pathologies, deep learning models show diagnostic performance matching or exceeding radiologist capabilities. Transfer learning approaches using established architectures (AlexNet, VGGNet, GoogLeNet, ResNet) achieve expert-level accuracy in otosclerosis detection [4]. In cochlear malformation diagnosis, AI systems outperform human specialists, demonstrating potential for clinical implementation [16]. Neural network analysis of chronic otitis media achieves 95.4% classification accuracy on temporal bone CT images, surpassing some radiologist performance benchmarks [17,18].

MRI-Based Diagnostics

AI applications in MRI-based diagnosis demonstrate consistent performance across various otologic conditions. For vestibular schwannoma (VS), a 2.5D neural network achieves 93.7% segmentation accuracy in tumor boundary delineation [19]. The technology additionally enables differentiation between solid and cystic tumor components while providing precise size measurements [20]. In Meniere's disease diagnosis, convolutional neural network (CNN) models reliably calculate endolymphatic hydrops ratios, with results showing strong correlation to expert evaluations [21].

Comparative analyses reveal AI systems match or exceed clinician performance across multiple diagnostic parameters. These systems achieve equivalent accuracy to specialists in detecting tympanic membrane pathologies [13], otosclerosis [4], and cochlear malformations [16], while demonstrating superior processing speeds - particularly in temporal bone CT segmentation tasks [5,15]. The technology additionally reduces inter-rater variability, providing standardized diagnostic outputs for complex cases [18,19]. (Table 1.)

Table 1. Diagnostic Applications of AI in Otology.

Diagnostic Area	AI Technology	Key Findings	Accuracy/Performance	Citation
Hearing Screening	Machine learning models	Classifies conductive/sensorineural/mixed hearing loss; aids in resource-limited settings.	–	[10, 11]
Tympanic Membrane (TM) Imaging	Deep learning (CNN-based)	Detects otitis media; outperforms physicians in pathology classification.	80–90% (vs. 75% clinicians)	[12]
	Diagnostic algorithm	Detects otitis media but cannot differentiate subtypes.	92.6%	[13]
	Subtyping model (Zafer et al.)	Differentiates AOM, OME, and CSOM.	–	[14]
Temporal Bone CT	W-Net method (Lv et al.)	Fully automatic segmentation; matches manual accuracy with faster processing.	–	[5, 15]
	Deep learning (AlexNet, VGGNet, etc.)	Diagnosis otosclerosis comparably to radiologists.	Matches radiologists	[4]
	CNN-based models	Identifies cochlear malformations better than physicians.	Superior to clinicians	[16]
	Neural networks	Classifies chronic otitis media and other temporal bone pathologies.	Up to 95.4%	[17, 18]
MRI Diagnostics	2.5D neural network	Segments vestibular schwannoma (VS) with high precision.	93.7%	[19]
	Deep learning model	Differentiates solid/cystic VS components and measures tumor size.	–	[20]
	CNN-based model	Calculates endolymphatic hydrops ratio in Ménière's disease; agrees with expert assessments.	High agreement	[21]

Advancing Hearing Loss Treatment with AI

AI in Hearing Aid Personalization

AI has significantly advanced hearing aid technology through automated, adaptive sound processing capabilities that reduce dependence on manual audiologist adjustments [22,23]. Deep neural networks (DNNs) now enable real-time analysis of acoustic environments, dynamically reducing background noise while enhancing speech clarity [24]. These systems automatically personalize amplification and noise reduction parameters based on individual user needs, substantially decreasing the number of required clinical visits for device optimization [23].

Cochlear Implant Optimization

AI enhances cochlear implantation outcomes through two principal mechanisms. First, electrode mapping algorithms predict optimal electrode placement patterns to maximize auditory nerve stimulation, resulting in improved speech perception outcomes. Second, AI-powered surgical planning systems analyze individual patient anatomy from preoperative imaging to guide surgical decision-making and improve procedural precision [25]. These applications demonstrate measurable improvements over conventional approaches while reducing subjective variability in clinical practice.

Predictive Therapeutics

AI applications show significant potential in managing sudden sensorineural hearing loss (SSHL). Deep belief networks (DBNs) demonstrate effectiveness in predicting treatment outcomes, providing valuable prognostic information to guide clinical decisions [26]. Furthermore, machine learning algorithms enable personalized therapeutic approaches by analyzing individual patient characteristics to optimize intervention strategies, including sound therapy and surgical options [27]. These technological advances offer data-driven alternatives to traditional standardized treatment protocols.

Vestibular Rehabilitation

AI demonstrates measurable benefits in vestibular rehabilitation through integration with virtual reality (VR) platforms and machine learning algorithms. Clinical studies document significant balance improvement in patients using AI-enhanced VR systems implementing Cawthorne-Cooksey exercises. The technology provides quantitative assessment of dysfunction severity through Support Vector Machine (SVM) analysis, enabling precise customization of rehabilitation protocols [28]. These applications show particular promise in standardizing and optimizing balance disorder treatment. (Table 2.)

Table 2. Therapeutic Applications of AI

Treatment Area	AI Technology Used	Key Functionality	Clinical Benefits	Citation
Hearing Aids	Deep Neural Networks (DNN)	Real-time environmental sound analysis and adaptive adjustment	Improved speech clarity in noise; reduced need for manual tuning	[22-24]
Cochlear Implants	Electrode mapping algorithms	Predictive modeling for optimal electrode placement	Better speech perception; reduced surgical complications	[25]
Sudden Hearing Loss	Deep Belief Networks (DBN)	Prognostic prediction based on clinical variables	Improved treatment decision-making for SSHL	[26]
Vestibular Rehabilitation	SVM + Virtual Reality	Automated impairment assessment and customized therapy programs	More effective balance therapy with objective progress tracking	[28]

Discussion

The progressive integration of artificial intelligence systems into daily audiological practice demonstrates significant advancements across multiple key domains, each presenting unique opportunities for care optimization as well as specific technological and implementational challenges. In the area of diagnostic applications, AI systems have proven to be highly effective, particularly in tasks requiring highly precise visual pattern recognition [10,11]. The capacity of advanced machine learning models to analyze tympanic membrane images with an accuracy of 80–90%, frequently surpassing the standard diagnostic performance of physicians (estimated at approximately 75%), emphatically underscores their immense potential as objective diagnostic adjuncts [12]. Nevertheless, the limitations and algorithmic errors observed, for instance, in the precise subtyping of otitis media, clearly demarcate the current boundaries of their technological capabilities. These indicate that while AI excels at the binary detection of pathology, it continues to face substantial difficulties with more nuanced clinical differentiation, where overlapping symptoms and atypical tympanic membrane presentations necessitate a broader medical context [13,14]. The medical discipline that is indisputably the fastest and most extensive in adapting new digital technologies is radiology [29]. Artificial intelligence solutions have found highly fertile ground in advanced imaging techniques utilized in audiology and neurotology, such as computed tomography (CT) and high-resolution magnetic resonance imaging (MRI) [30]. In the diagnostic workup of otological disorders, computed tomography of the temporal bone frequently constitutes an absolutely indispensable component of the examination, determining subsequent therapeutic management [31]. The anatomical complexity of the middle and inner ear means that in this domain, AI possesses immense potential to support physicians in their daily practice. Recent scientific reports unequivocally indicate that appropriately trained algorithms can significantly streamline the diagnostic process, offering not only faster generation of results but, above all, minimizing the risk of oversights and errors resulting from perceptual fatigue compared to human specialists [32]. The development of fully automated segmentation algorithms that can match the accuracy of tedious manual segmentation, while simultaneously achieving a drastic reduction in data processing time, addresses a long-standing clinical demand [15]. Until recently, the process of precise segmentation of individual structures was performed exclusively manually, which consumed disproportionate amounts of time; therefore, intensive research into partial or complete automation of this stage has long been conducted within the scientific community [33]. These advancements are particularly invaluable in the assessment of exceptionally complex anatomical conditions, where AI demonstrates significantly higher efficacy in the flawless identification of subtle cochlear malformations and early otosclerotic foci compared to traditional, visual methods of radiological assessment [4,16]. The impressive accuracy of 93.7% achieved in the automated segmentation of vestibular schwannomas further highlights the utility of AI in the analysis of cerebellopontine angle tumor growth, although it must be noted that the absolute requirement for highly standardized, high-quality training datasets may continue to restrict the broader application of these tools in smaller centers in the near future [19]. In therapeutic applications, the revolutionary transition from static devices to fully adaptive hearing aid technologies, based on deep neural networks, represents a fundamental and arguably the most significant paradigm shift in hearing rehabilitation in decades [23,24]. This rapid evolution—from simple parameter programming by a prosthetist to the real-time, autonomous environmental adaptation of the algorithm—effectively eliminates the most severe limitation of traditional hearing aids: their inability to dynamically and instantaneously respond to rapidly changing and highly complex acoustic environments [22]. Similarly, AI-assisted software for mapping sound processors in cochlear implants can propose significantly more precise and personalized patterns of neural tissue stimulation, directly translating into the potential optimization of speech understanding and perception outcomes in implanted patients [25]. The application of deep belief networks to predict treatment outcomes in cases of sudden sensorineural hearing loss introduces an exceptionally valuable prognostic tool to clinicians. Nevertheless, physicians must exercise considerable caution to avoid falling into excessive, uncritical reliance on algorithmic predictions, especially when the decision-making mechanisms underlying a specific outcome remain opaque or impossible to explain to them (the so-called 'black box' problem) [26]. Researchers hold similar concerns regarding modern vestibular rehabilitation systems utilizing virtual reality (VR) environments. Although the preliminary results of these interventions are highly optimistic, their long-term clinical utility and the durability of effects require rigorous validation through broader, multicenter studies on highly diverse populations of patients with various types of balance disorders [28]. However, several critical systemic challenges emerge from the rapid development of these technologies. First, the noticeable variability in the performance of analytical models depending on diverse clinical settings unequivocally suggests that current AI systems may lack the generalizability and robustness essential for their global, safe implementation [12,16].

Second, the very high computational and infrastructural demands necessary to support many of the described applications may generate new disparities in access to modern medical care, creating a form of digital exclusion for healthcare facilities located in resource-limited environments [15,19]. Third, the physical integration of these advanced technologies into existing, often outdated hospital software and the rigid workflows of clinical staff poses tremendous practical challenges, the resolution of which requires an entirely different approach than merely striving to improve the technical efficacy metrics of the algorithms themselves [22,25].

Challenges and Ethical Considerations

Despite its immense and transformative potential, the widespread integration of artificial intelligence into audiological and otorhinolaryngological practice is not without profound systemic and methodological challenges. Foremost among these is the critical dependency of machine learning algorithms—particularly deep neural networks—on the availability of massive, meticulously labelled, and highly diverse training datasets. The current scarcity of such comprehensive data repositories constitutes a significant bottleneck, frequently limiting the scope, accuracy, and generalizability of contemporary research. Furthermore, AI-generated prognostic and diagnostic predictions in medicine must undergo rigorous, prospective clinical validation across heterogeneous patient populations. This step is indispensable, as inherent algorithmic uncertainty, data bias, and other methodological disparities can substantially compromise clinical reliability and potentially lead to erroneous therapeutic decisions [34].

Another paramount challenge lies in ensuring the transparency and interpretability of complex AI algorithms, often referred to as the "black box" dilemma. For these advanced computational models to be safely and effectively incorporated into routine practice, clinicians must have a clear understanding of the underlying parameters and decision-making pathways that lead to specific diagnostic conclusions. Establishing this algorithmic transparency is foundational to building clinical trust and ensuring that these powerful tools are utilized appropriately as adjuncts rather than replacements for clinical judgment. Consequently, the gradual, phased implementation of AI solutions, firmly rooted in evidence-based best practices and supported by continuous, multi-directional feedback mechanisms between end-users and developers, is essential to systematically mitigate the prevailing skepticism and hesitance among healthcare professionals [35].

In addition to technical hurdles, the extraordinarily dynamic evolution of artificial intelligence technologies necessitates a paradigm shift in medical education, requiring otorhinolaryngologists to engage in continuous, lifelong skill acquisition. The development and provision of specialized educational initiatives—encompassing interactive workshops, advanced seminars, and comprehensive online curricula focused on specific AI applications, such as algorithms for the automated diagnosis of otitis media or the objective acoustic assessment of voice disorders—are of crucial importance. To bridge the gap between technological innovation and clinical application, further robust research and the formulation of standardized, professional guidelines are urgently needed. Ultimately, hands-on, practical training dedicated to the seamless integration of AI tools into daily clinical workflows will empower clinicians to leverage these technologies safely, thereby maintaining and elevating the highest standards of patient-centric care [36].

Finally, ethical considerations occupy a central and highly sensitive role in the deployment of AI in healthcare, particularly concerning the stringency of data privacy and the complexities of obtaining truly informed patient consent. Audiological and otoneurological data—encompassing highly sensitive personal health information such as audiometric test results, vocal biomarkers, and high-resolution radiological imaging—must be encrypted, stored, and processed within exceptionally secure, compliant infrastructures to preserve absolute patient confidentiality. Addressing these multifaceted ethical imperatives requires an unprecedented level of interdisciplinary collaboration. It is imperative that healthcare policymakers, clinical ethicists, and software developers work synergistically to establish robust, standardized, and legally binding guidelines governing the ethical development, deployment, and continuous auditing of artificial intelligence systems in audiology [37].

Conclusions

Artificial intelligence stands on the precipice of playing a profoundly transformative role within the field of audiology and otorhinolaryngology. By introducing highly sophisticated computational tools and predictive models, AI holds the capacity to significantly enhance diagnostic precision, dynamically optimize the performance parameters of advanced hearing rehabilitative devices, and ultimately elevate overall long-term patient clinical outcomes. As machine learning architectures and deep learning technologies continue to evolve at an unprecedented pace, fostering robust interdisciplinary collaboration will become increasingly indispensable. The synergistic integration of clinical expertise from audiologists and medical practitioners with the technical proficiency of data scientists, software engineers, and bioethicists will be crucial to fully, safely, and effectively harness the immense potential of AI within evidence-based clinical practice. Furthermore, by systematically addressing the current infrastructural, computational, and methodological challenges, and by strictly prioritizing paramount ethical considerations—particularly regarding data security, patient privacy, and algorithmic transparency—artificial intelligence can pave the way for a monumental paradigm shift in healthcare. This technological evolution promises a future where the diagnosis, continuous management, and targeted treatment of complex hearing and balance disorders are executed with unparalleled clinical precision, therapeutic efficacy, and operational efficiency.

Conflict of Interest Statement: The authors declare no conflict of interest.

Funding Sources: There are no sources of funding to declare.

REFERENCES

1. Koyama, H., Kashio, A., & Yamasoba, T. (2024). Application of artificial intelligence in otology: Past, present, and future. *Journal of Clinical Medicine*, 13(24), 7577. <https://doi.org/10.3390/jcm13247577>
2. Frosolini, A., Franz, L., Caragli, V., Genovese, E., de Filippis, C., & Marioni, G. (2024). Artificial intelligence in audiology: A scoping review of current applications and future directions. *Sensors*, 24(22), 7126. <https://doi.org/10.3390/s24227126>
3. Wang, S., Mo, C., Chen, Y., Dai, X., Wang, H., & Shen, X. (2024). Exploring the performance of ChatGPT-4 in the Taiwan Audiologist Qualification Examination: Preliminary observational study highlighting the potential of AI chatbots in hearing care. *JMIR Medical Education*, 10, e55595. <https://doi.org/10.2196/55595>
4. Fujima, N., Andreu-Arasa, V. C., Onoue, K., Weber, P. C., Hubbell, R. D., Setty, B. N., & Sakai, O. (2021). Utility of deep learning for the diagnosis of otosclerosis on temporal bone CT. *European Radiology*, 31(7), 5206–5211. <https://doi.org/10.1007/s00330-020-07568-0>
5. Lv, Y., Ke, J., Xu, Y., Shen, Y., Wang, J., & Wang, J. (2021). Automatic segmentation of temporal bone structures from clinical conventional CT using a CNN approach. *The International Journal of Medical Robotics and Computer Assisted Surgery*, 17(2), e2229. <https://doi.org/10.1002/rcs.2229>
6. Rodrigo, H., Beukes, E. W., Andersson, G., & Manchaiah, V. (2021). Exploratory data mining techniques (decision tree models) for examining the impact of internet-based cognitive behavioral therapy for tinnitus: Machine learning approach. *Journal of Medical Internet Research*, 23(11), e28999. <https://doi.org/10.2196/28999>
7. Mondol, S. I. M. M. R., Kim, H. J., Kim, K. S., & Lee, S. (2022). Machine learning-based hearing aid fitting personalization using clinical fitting data. *Journal of Healthcare Engineering*, 2022, 1667672. <https://doi.org/10.1155/2022/1667672>
8. Zhang, J., & Zhang, Z. M. (2023). Ethics and governance of trustworthy medical artificial intelligence. *BMC Medical Informatics and Decision Making*, 23(1), 7. <https://doi.org/10.1186/s12911-023-02103-9>
9. Bur, A. M., Shew, M., & New, J. (2019). Artificial intelligence for the otolaryngologist: A state of the art review. *Otolaryngology–Head and Neck Surgery*, 160(4), 603–611. <https://doi.org/10.1177/0194599819827507>
10. Wang, Y., Yao, X., Wang, D., Ye, C., & Xu, L. (2024). A machine learning screening model for identifying the risk of high-frequency hearing impairment in a general population. *BMC Public Health*, 24(1), 1160. <https://doi.org/10.1186/s12889-024-18636-1>

11. Gathman, T. J., Choi, J. S., Vasdev, R. M. S., Schoepfoerster, J. A., & Adams, M. E. (2023). Machine learning prediction of objective hearing loss with demographics, clinical factors, and subjective hearing status. *Otolaryngology–Head and Neck Surgery*, 169(3), 504–513. <https://doi.org/10.1002/ohn.288>
12. Cao, Z., Chen, F., Grais, E. M., Yue, F., Cai, Y., Swanepoel, W., & Zhao, F. (2023). Machine learning in diagnosing middle ear disorders using tympanic membrane images: A meta-analysis. *The Laryngoscope*, 133(4), 732–741. <https://doi.org/10.1002/lary.30291>
13. Sundgaard, J. V., Bray, P., Laugesen, S., Harte, J., Kamide, Y., Tanaka, C., Christensen, A. N., & Paulsen, R. R. (2022). A deep learning approach for detecting otitis media from wideband tympanometry measurements. *IEEE Journal of Biomedical and Health Informatics*, 26(7), 2974–2982. <https://doi.org/10.1109/JBHI.2022.3159263>
14. Cömert, Z. (2019). Fusing fine-tuned deep features for recognizing different tympanic membranes. *Biocybernetics and Biomedical Engineering*, 40(1), 40–51. <https://doi.org/10.1016/j.bbe.2019.11.001>
15. Xianfen, D., Siping, C., Changhong, L., & Yuanmei, W. (2005). 3D semi-automatic segmentation of the cochlea and inner ear. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2005*, 6285–6288. <https://doi.org/10.1109/IEMBS.2005.1615934>
16. Li, Z., Zhou, L., Bin, X., Tan, S., Tan, Z., & Tang, A. (2024). Utility of deep learning for the diagnosis of cochlear malformation on temporal bone CT. *Japanese Journal of Radiology*, 42(3), 261–267. <https://doi.org/10.1007/s11604-023-01494-z>
17. Wang, Y. M., Li, Y., Cheng, Y. S., He, Z. Y., Yang, J. M., Xu, J. H., Chi, Z. C., Chi, F. L., & Ren, D. D. (2020). Deep learning in automated region proposal and diagnosis of chronic otitis media based on computed tomography. *Ear and Hearing*, 41(3), 669–677. <https://doi.org/10.1097/AUD.0000000000000794>
18. Eroğlu, O., Eroğlu, Y., Yıldırım, M., Karlıdag, T., Çınar, A., Akyiğit, A., Kaygusuz, İ., Yıldırım, H., Keleş, E., & Yalçın, Ş. (2022). Is it useful to use computerized tomography image-based artificial intelligence modelling in the differential diagnosis of chronic otitis media with and without cholesteatoma? *American Journal of Otolaryngology*, 43(3), 103395. <https://doi.org/10.1016/j.amjoto.2022.103395>
19. Shapey, J., Wang, G., Dorent, R., Dimitriadis, A., Li, W., Paddick, I., Kitchen, N., Bisdas, S., Saeed, S. R., Ourselin, S., Bradford, R., & Vercauteren, T. (2019). An artificial intelligence framework for automatic segmentation and volumetry of vestibular schwannomas from contrast-enhanced T1-weighted and high-resolution T2-weighted MRI. *Journal of Neurosurgery*, 134(1), 171–179. <https://doi.org/10.3171/2019.9.JNS191949>
20. Neve, O. M., Chen, Y., Tao, Q., Romeijn, S. R., de Boer, N. P., Grootjans, W., Kruit, M. C., Lelieveldt, B. P. F., Jansen, J. C., Hensen, E. F., Verbist, B. M., & Staring, M. (2022). Fully automated 3D vestibular schwannoma segmentation with and without gadolinium-based contrast material: A multicenter, multivendor study. *Radiology: Artificial Intelligence*, 4(4), e210300. <https://doi.org/10.1148/ryai.210300>
21. Cho, Y. S., Cho, K., Park, C. J., Chung, M. J., Kim, J. H., Kim, K., Kim, Y. K., Kim, H. J., Ko, J. W., Cho, B. H., & Chung, W. H. (2020). Automated measurement of hydrops ratio from MRI in patients with Ménière's disease using CNN-based segmentation. *Scientific Reports*, 10(1), 7003. <https://doi.org/10.1038/s41598-020-63887-8>
22. Convery, E., Keidser, G., Seeto, M., Yeend, I., & Freeston, K. (2015). Factors affecting reliability and validity of self-directed automatic in situ audiometry: Implications for self-fitting hearing aids. *Journal of the American Academy of Audiology*, 26(1), 5–18. <https://doi.org/10.3766/jaaa.26.1.2>
23. Ghosh Moulic, A., Gaurkar, S. S., & Deshmukh, P. T. (2024). Artificial intelligence in otology, rhinology, and laryngology: A narrative review of its current and evolving picture. *Cureus*, 16(8), e66036. <https://doi.org/10.7759/cureus.66036>
24. Saoji, A. A., Sheikh, B. A., Bertsch, N. J., Goulson, K. R., Graham, M. K., McDonald, E. A., Bross, A. E., Vaisberg, J. M., Kühnel, V., Voss, S. C., Qian, J., Hogan, C. H., & DeJong, M. D. (2024). How does deep neural network-based noise reduction in hearing aids impact cochlear implant candidacy? *Audiology Research*, 14(6), 1114–1125. <https://doi.org/10.3390/audiolres14060092>
25. Essaid, B., Kheddar, H., Batel, N., Chowdhury, M. E. H., & Lakas, A. (2024). Artificial intelligence for cochlear implants: Review of strategies, challenges, and perspectives. *IEEE Access*, 12, 119015–119038. <https://doi.org/10.1109/ACCESS.2024.3429524>
26. Bing, D., Ying, J., Miao, J., Lan, L., Wang, D., Zhao, L., Yin, Z., Yu, L., Guan, J., & Wang, Q. (2018). Predicting the hearing outcome in sudden sensorineural hearing loss via machine learning models. *Clinical Otolaryngology*, 43(3), 868–874. <https://doi.org/10.1111/coa.13068>

27. Zhang, G., Chen, R., Ghorbani, H., Li, W., Minasyan, A., Huang, Y., Lin, S., & Shao, M. (2025). Artificial intelligence-enabled innovations in cochlear implant technology: Advancing auditory prosthetics for hearing restoration. *Bioengineering & Translational Medicine*. Advance online publication. <https://doi.org/10.1002/btm2.10752>
28. Yeh, S. C., Huang, M. C., Wang, P. C., Fang, T. Y., Su, M. C., Tsai, P. Y., & Rizzo, A. (2014). Machine learning-based assessment tool for imbalance and vestibular dysfunction with virtual reality rehabilitation system. *Computer Methods and Programs in Biomedicine*, 116(3), 311–318. <https://doi.org/10.1016/j.cmpb.2014.04.014>
29. Ankush, A. (2025). VoxRad: Building an open-source locally-hosted radiology reporting system. *Clinical Imaging*, 119, 110414. <https://doi.org/10.1016/j.clinimag.2025.110414>
30. Kujawa, A., Dorent, R., Connor, S., Thomson, S., Ivory, M., Vahedi, A., Guilhem, E., Wijethilake, N., Bradford, R., Kitchen, N., Bisdas, S., Ourselin, S., Vercauteren, T., & Shapey, J. (2024). Deep learning for automatic segmentation of vestibular schwannoma: A retrospective study from multi-center routine MRI. *Frontiers in Computational Neuroscience*, 18, 1365727. <https://doi.org/10.3389/fncom.2024.1365727>
31. Matari, N. H., Mingomataj, E., Famuyide, A., Nguyen, P., Lalwani, A. K., & Garcon, E. (2025). Pöschl reformations created from high-resolution noncontrast enhanced CT head exams can be used to detect and classify superior semicircular canal abnormalities. *Otology & Neurotology*, 46(2), 176–182. <https://doi.org/10.1097/MAO.0000000000004395>
32. Spinos, D., Martinos, A., Petsiou, D. P., Mistry, N., & Garas, G. (2025). Artificial intelligence in temporal bone imaging: A systematic review. *The Laryngoscope*, 135(3), 973–981. <https://doi.org/10.1002/lary.31809>
33. Xianfen, D., Siping, C., Changhong, L., & Yuanmei, W. (2005). 3D semi-automatic segmentation of the cochlea and inner ear. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society, 2005*, 6285–6288. <https://doi.org/10.1109/IEMBS.2005.1615934>
34. Tama, B. A., Kim, D. H., Kim, G., Kim, S. W., & Lee, S. (2020). Recent advances in the application of artificial intelligence in otorhinolaryngology-head and neck surgery. *Clinical and Experimental Otorhinolaryngology*, 13(4), 326–339. <https://doi.org/10.21053/ceo.2020.00654>
35. Huang, Z., George, M. M., Tan, Y. R., Natarajan, K., Devasagayam, E., Tay, E., Manesh, A., Varghese, G. M., Abraham, O. C., Zachariah, A., Yap, P., Lall, D., & Chow, A. (2023). Are physicians ready for precision antibiotic prescribing? A qualitative analysis of the acceptance of artificial intelligence-enabled clinical decision support systems in India and Singapore. *Journal of Global Antimicrobial Resistance*, 35, 76–85. <https://doi.org/10.1016/j.jgar.2023.08.016>
36. Irfan, B. (2024). Beyond the scope: Advancing otolaryngology with artificial intelligence integration. *Cureus*, 16(2), e54248. <https://doi.org/10.7759/cureus.54248>
37. Lechien, J. R. (2024). Generative artificial intelligence in otolaryngology-head and neck surgery editorial: Be an actor of the future or follower. *European Archives of Oto-Rhino-Laryngology*, 281(4), 2051–2053. <https://doi.org/10.1007/s00405-024-08579-z>