



International Journal of Innovative Technologies in Social Science

e-ISSN: 2544-9435

Operating Publisher
SciFormat Publishing Inc.
ISNI: 0000 0005 1449 8214

2734 17 Avenue SW,
Calgary, Alberta, T3E0A7,
Canada
+15878858911
editorial-office@sciformat.ca

ARTICLE TITLE	LIFE-THREATENING INFECTIOUS CONSEQUENCES OF NEGLECTED ORAL CAVITY DISEASES: A COMPREHENSIVE REVIEW
----------------------	--

DOI	https://doi.org/10.31435/ijitss.3(51).2026.6720
------------	---

RECEIVED	18 July 2026
-----------------	--------------

ACCEPTED	17 September 2026
-----------------	-------------------

PUBLISHED	25 September 2026
------------------	-------------------

LICENSE



The article is licensed under a **Creative Commons Attribution 4.0 International License**.

© The author(s) 2026.

This article is published as open access under the Creative Commons Attribution 4.0 International License (CC BY 4.0), allowing the author to retain copyright. The CC BY 4.0 License permits the content to be copied, adapted, displayed, distributed, republished, or reused for any purpose, including adaptation and commercial use, as long as proper attribution is provided.

LIFE-THREATENING INFECTIOUS CONSEQUENCES OF NEGLECTED ORAL CAVITY DISEASES: A COMPREHENSIVE REVIEW

Wojciech Gawęda (Corresponding Author, Email: gaweda.wojtek@gmail.com)
Medical University of Lublin, Aleje Raławickie 1, 20-059 Lublin, Poland
ORCID ID: 0009-0007-2461-8584

Wiktoria Łobodzińska
Medical University of Lublin, Aleje Raławickie 1, 20-059 Lublin, Poland
ORCID ID: 0009-0003-1011-7915

Maciej Ficek
Medical University of Lublin, Aleje Raławickie 1, 20-059 Lublin, Poland
ORCID ID: 0009-0003-7629-448X

Justyna Adamczyk
Medical University of Lublin, Aleje Raławickie 1, 20-059 Lublin, Poland
ORCID ID: 0009-0008-6548-8998

Kinga Głodek
Medical University of Lublin, Aleje Raławickie 1, 20-059 Lublin, Poland
ORCID ID: 0009-0006-8126-0702

Martyna Kowalczyk
Wrocław Medical University, Ludwik Pasteur 1, 50-367 Wrocław, Poland
ORCID ID: 0009-0009-5505-2776

ABSTRACT

Background: Untreated dental caries, pulpal gangrene, apical periodontitis, and impacted molar pericoronitis remain prevalent chronic conditions that frequently breach local biological containment when professional intervention is withheld or delayed.

Objectives: To evaluate the direct pathophysiological routes, microbiological shifts, and clinical spectrum of life-threatening regional and systemic infections specifically arising from untreated oral diseases.

Methods: A narrative synthesis of peer-reviewed clinical studies, prospective trials, and systematic reviews indexed in PubMed addressing severe odontogenic deep neck infections, mediastinal extension, intracranial spreading, and necrotizing fasciitis was conducted.

Results: Odontogenic origins account for 34.2% to 49% of all diagnosed deep neck infections. Cortical erosion inferior to the mylohyoid line precipitates Ludwig's angina, leading to tongue elevation and precipitous airway asphyxia. Uncontrolled extension through the pharyngeal and retrovisceral 'danger space' (Space 4) causes descending necrotizing mediastinitis, carrying a mortality rate of 20–40%. Craniomaxillofacial necrotizing fasciitis displays a striking 30.3% mortality rate in diabetic cohorts due to severe microvascular thrombosis. Retrograde propagation via valveless ophthalmic and facial venous plexuses precipitates cavernous sinus thrombosis, orbital cellulitis, and brain abscesses. Microbiological evaluation demonstrates penicillin-resistant strains in 54% of severe cases driven by beta-lactamase-producing anaerobes, rendering delayed antimicrobial monotherapy ineffective without surgical intervention.

Conclusions: Abandoning routine and curative dental care directly triggers catastrophic surgical emergencies. Early identification of red-flag symptoms (trismus, dysphagia, stridor), broad-spectrum parenteral coverage, and prompt surgical eradication of the causative dental source are imperative to prevent fatal outcomes and long-term musculoskeletal deformity.

KEYWORDS

Odontogenic Infections, Dental Neglect, Ludwig's Angina, Descending Necrotizing Mediastinitis, Necrotizing Fasciitis, Cavernous Sinus Thrombosis, Deep Neck Space Infection

CITATION

Gawęda, W., Łobodzińska, W., Ficek, M., Adamczyk, J., Głodek, K., & Kowalczyk, M. (2026). Life-threatening infectious consequences of neglected oral cavity diseases: A comprehensive review. *International Journal of Innovative Technologies in Social Science*, 3(51). [https://doi.org/10.31435/ijitss.3\(51\).2026.6720](https://doi.org/10.31435/ijitss.3(51).2026.6720)

COPYRIGHT

© The author(s) 2026. This article is published as open access under the **Creative Commons Attribution 4.0 International License (CC BY 4.0)**, allowing the author to retain copyright. The CC BY 4.0 License permits the content to be copied, adapted, displayed, distributed, republished, or reused for any purpose, including adaptation and commercial use, as long as proper attribution is provided.

The Natural Progression of Untreated Oral Cavity Diseases

Dental caries and chronic periodontal destruction represent two of the most ubiquitous chronic pathologies in human populations, yet their potential to trigger catastrophic, systemic crises is frequently underestimated when patients avoid or delay professional dental care [1,7]. The progression from an initial, asymptomatic enamel breakdown into an invasive head and neck crisis represents a continuous pathophysiological continuum. When debridement, endodontic therapy, or surgical extraction is omitted, bacterial enzymes continuously demineralize dental hard tissues, penetrating the dentinal tubules and invading the pulp chamber [3,7]. This uninhibited bacterial colonization incites severe inflammatory hyperemia, which rapidly leads to strangulation of the pulp microvasculature at the narrow apical foramen, culminating in complete pulp gangrene and liquefactive necrosis [3,7]. Deprived of host immune defenses, the necrotic root canal architecture transforms into an uninhibited incubation reservoir for highly virulent anaerobic polymicrobial communities [3].

As microbial replication intensifies within the root canal, septic exudate escapes through the apical foramen into the surrounding periodontal ligament, instigating periapical periodontitis [3,7]. Osteoclastic activation driven by bacterial endotoxins resorbs alveolar cancellous bone, gradually expanding the periapical lesion into an acute alveolar abscess [3]. With growing purulent accumulation within the enclosed gnathic bone, hydrostatic pressure rises until the exudate perforates the cortical plate along the path of least resistance [1,7]. The anatomical position of this osseous perforation relative to specific muscular attachments dictates whether the infection remains superficial or bursts into deep anatomical spaces [1,3,7]. In the mandible, the insertion of the mylohyoid muscle forms the definitive anatomical watershed: infections from incisors, canines, and premolars breach the cortical plate superior to the mylohyoid line, dumping pus into the sublingual space, whereas the root apices of the second and third mandibular molars lie inferior to the mylohyoid attachment, directing uncontrolled suppuration directly into the submandibular and submental compartments [1,3,7]. Similarly, pericoronitis arising from semi-impacted lower third molars represents a notoriously dangerous portal of entry; the accumulation of necrotic debris under an inflamed operculum grants virulent pathogens direct access into the masticator spaces (submasseteric, pterygomandibular, and temporal spaces) and lateral pharyngeal fascial conduits [1,3,6].

Direct Life-Threatening Regional and Systemic Complications

The failure to address primary oral pathologies removes all biological barriers, allowing localized dental abscesses to breach the deep cervical fascia. Data across maxillofacial surgery units demonstrate that untreated odontogenic foci account for 34.2% to 49% of all diagnosed deep neck infections (DNIs), which represent acute, life-threatening surgical emergencies [1,2,4]. The resulting clinical manifestations depend directly on the fascial pathways breached by the untreated dental pathogen reservoir [1,6].

a) Airway Compromise and Ludwig's Angina

A catastrophic consequence of untreated lower molar apical periodontitis or pericoronitis is the rapid establishment of Ludwig's angina [1,4,6]. This clinical entity consists of an aggressive, rapidly expanding, brawny cellulitis that engulfs the submandibular, sublingual, and submental spaces bilaterally [2,4,6]. Unlike typical fluctuant abscesses, Ludwig's angina is characterized in its early stages by dense, board-like induration without localized pus cavities, rendering spontaneous drainage impossible [4,6]. As massive edema accumulates within the unyielding confines of the floor of the mouth, the tongue is forcefully elevated and

pushed backward against the posterior pharyngeal wall and soft palate [1,2,4,6]. This posterior mechanical displacement causes acute, precipitous upper airway occlusion [1,4]. Patients exhibit profound trismus resulting from the irritation of the masticator space musculature, severe odynophagia, inability to swallow their own saliva, a distinctive muffled 'hot potato' voice, and tachypnea, culminating in sudden asphyxiation if the airway is not urgently stabilized via awake fiberoptic intubation or emergency surgical tracheostomy [1,2,4,6].

b) Descending Necrotizing Mediastinitis (DNM)

When a neglected lower molar or pericoronal infection penetrates through the posterior boundary of the mylohyoid muscle, it directly enters the lateral pharyngeal (parapharyngeal) space [1,2,6]. From this space, purulence tracks unchecked into the retropharyngeal space and the anatomically critical 'danger space' (Space 4), bounded anteriorly by the alar fascia and posteriorly by the prevertebral fascia [2,6]. Because the danger space consists of loose, areolar connective tissue that offers zero anatomical resistance, it provides an open conduit extending from the base of the skull directly into the posterior mediastinum at the level of the diaphragm [2,6,7]. Gravity, combined with the negative intrathoracic pressures generated during normal inspiratory cycles and the continuous pumping motion of swallowing, actively draws purulent exudate down through the neck into the thoracic cavity [2,6]. The patient deteriorates rapidly, presenting with retrosternal chest pain, dyspnea, pleural empyema, purulent pericarditis, and refractory septic shock [2,6,7]. Despite modern imaging and intensive thoracic surgical interventions, descending necrotizing mediastinitis secondary to untreated dental disease continues to carry an alarming mortality rate exceeding 20% to 40% [2,6,7].

c) Craniomaxillofacial and Cervical Necrotizing Fasciitis

Another severe outcome of neglected oral suppurative disease is odontogenic necrotizing fasciitis (ONF), an aggressive bacterial infection marked by widespread necrosis of subcutaneous tissue and superficial and deep cervical fascia [5,6,7]. Bacterial proliferation and intense enzymatic release prompt widespread thrombosis of nutrient microvessels coursing through fascial planes [2,5]. As cutaneous blood supply is severed, the overlying skin rapidly transitions from erythema and tense edema to dusky, violaceous discoloration, bullae formation, and frank gangrene [5,6]. A cardinal diagnostic clue is severe pain disproportionate to visible swelling, followed quickly by cutaneous anesthesia caused by ischemic destruction of superficial sensory nerves, accompanied by subcutaneous crepitus from gas-producing anaerobes [2,5,6]. Systematic reviews document an overall mortality of 9.8%, which escalates dramatically to 30.3% in patients with comorbid diabetes mellitus (odds ratio: 9.1, $p = 0.0001$) [5]. In diabetic hosts, impaired leukocyte chemotaxis, defective phagocytosis, and microvascular baseline damage permit unchecked bacterial dissection along cervical sheaths, frequently leading to carotid artery erosion, fatal jugular vein rupture, systemic toxemia, and severe cosmetic mutilation requiring extensive reconstructive surgery [2,5,6].

d) Craniomaxillofacial and Intracranial Complications

Neglecting teeth in either the maxilla or mandible can provoke intracranial complications via contiguous, retrograde venous, or hematogenous routes [6,7,8]. In the upper jaw, untreated caries or apical abscesses of premolars and molars erode the paper-thin maxillary sinus floor, causing chronic odontogenic maxillary sinusitis [6,7]. If ignored, purulence traverses the infraorbital canal, pterygopalatine fossa, and inferior orbital fissure, invading the orbit to cause orbital cellulitis, subperiosteal abscesses, chemosis, proptosis, and permanent blindness due to optic nerve compression or ischemic necrosis [6,7]. Simultaneously, retrograde propagation of septic thrombi through the valveless facial, angular, and superior ophthalmic veins—or posteriorly via the pterygoid venous plexus—allows pathogens to seed the cavernous sinus, producing septic cavernous sinus thrombosis [6,7]. Patients present with severe periorbital edema, external ophthalmoplegia resulting from paralysis of cranial nerves III, IV, and VI, loss of corneal reflex and forehead sensation (cranial nerve V1 and V2), and high fatality [6,7]. Furthermore, neglected oral chronic septic foci are established etiologic reservoirs for intracranial subdural empyemas, epidural abscesses, and cerebral abscesses predominantly situated in the frontal and temporal lobes [6,7,8]. Patients frequently experience an insidious onset of persistent headache, cognitive decline, seizures, progressive focal neurological deficits, and coma [Ref. 8].

e) Vascular Erosion, Internal Jugular Thrombophlebitis, and Sepsis

When purulence from untreated lower molars invades the parapharyngeal space, it surrounds the carotid sheath, threatening the major neurovascular structures of the neck [2,4,6]. Suppuration can cause septic thrombophlebitis of the internal jugular vein (a severe clinical presentation mirroring Lemierre's syndrome) [2,4,6]. Fragile septic fragments continuously detach into the bloodstream, producing metastatic septic emboli throughout the pulmonary vasculature, resulting in cavitating lung abscesses, empyema, and respiratory failure [2,4,6]. If host containment mechanisms fail, the relentless shower of bacteria and endotoxins into the circulation triggers systemic inflammatory response syndrome (SIRS), refractory septic shock, disseminated intravascular coagulation (DIC), and multiple organ dysfunction syndrome (MODS) [2,4,6,7].

Microbiological Evolution and Antibiotic Resistance in Neglected Infections

The clinical behavior of spreading dental infections is intrinsically tied to the polymicrobial ecology of the oral cavity and its evolution over time [3,4,7]. In the early phase of pulpal infection, the microflora is characterized predominantly by facultative aerobic streptococci, including *Streptococcus viridans*, *S. anginosus*, *S. constellatus*, and *S. intermedius* [3,4,5,7]. These early colonizers consume localized oxygen and generate an acidic tissue microenvironment, dramatically lowering the oxidation-reduction potential of the affected tissues [3]. As tissue necrosis advances in the absence of treatment, strict obligate anaerobes begin to dominate the expanding infection [3,4,5,6]. Species such as *Prevotella*, *Porphyromonas*, *Fusobacterium nucleatum*, *Peptostreptococcus*, and *Bacteroides* proliferate within the ischemic, deoxygenated fascial planes [3,4,5,6].

These anaerobic species produce aggressive virulence factors, notably hyaluronidase, chondroitin sulfatase, collagenase, and heparinase, which actively dissolve intercellular ground substance, destroy connective tissue sheaths, and induce microvascular thrombosis [3,5,7]. This enzymatic breakdown liquefies fascial barriers, permitting the rapid anatomical dissection described in deep neck space infections [3,7]. Importantly, the delay in seeking dental treatment frequently coincides with repeated, inappropriate courses of self-administered or empiric low-dose oral antibiotics, driving antimicrobial resistance [3]. Prospective microbiological evaluations have demonstrated that penicillin-resistant bacterial strains occur in 54% of hospitalized patients with severe odontogenic infections, with an overall penicillin clinical failure rate of approximately 21% [3]. This resistance is driven predominantly by beta-lactamase production among oral anaerobic species, including *Prevotella* and *Fusobacterium* [3]. Consequently, delayed pharmacological treatment with penicillin monotherapy is largely ineffective; overcoming these advanced infections requires broad-spectrum parenteral coverage (such as beta-lactam/beta-lactamase inhibitor combinations or carbapenems) combined with urgent, definitive surgical decompression and physical removal of the offending tooth [2,3,5].

Structural and Osseous Breakdown: Osteomyelitis and Permanent Deformity

The consequences of neglecting primary oral disease are not limited to acute fascial emergencies; prolonged chronic suppuration induces profound irreversible destruction of the gnathic skeleton and adjacent structures [1,6,7]. When a patient neglects chronic apical or marginal periodontitis for months or years, continuous purulent production within the alveolar bone disrupts the intraosseous vascular network and strips the periosteum away from the mandibular or maxillary cortex [1,6]. This loss of blood supply leads to diffuse chronic suppurative osteomyelitis of the jaws [1,6]. The necrotic bone separates from living tissue, forming large cortical sequestra that perpetuate active infection, stimulate painful fistulation through the facial skin, and cause spontaneous pathological jaw fractures [1,6].

In posterior mandibular molar infections, chronic osteolytic processes can track along the mandibular ramus directly to the temporomandibular joint (TMJ) [1,6]. The resulting septic arthritis and destructive condylar osteonecrosis dissolve the articular surfaces, culminating in severe fibrous or bony ankylosis of the jaw [1,6]. Patients are left with permanent mandibular hypomobility, severe restrictions in oral opening, and chronic masticatory impairment [1,6]. Furthermore, the radical surgical decompression, wide fasciotomies, and serial surgical debridements required to rescue patients from neglected multi-space infections inevitably result in substantial scar contractures, facial disfigurement, and irreversible transection of peripheral nerves—most notably the marginal mandibular branch of the facial nerve [2,5,7]. Thus, the initial omission of basic dental treatment leaves patients burdened with lifelong functional, anatomical, and psychological disabilities [2,5].

Integrative Synthesis of Neglected Dental Entities and Lethal Trajectories

An integrative analysis of untreated dental pathologies demonstrates how specific primary odontogenic conditions map directly to catastrophic, life-threatening clinical entities based on anatomical topography and microbial virulence [1,2,6,7]. In the lower jaw, deep dental caries and pulpal necrosis of the second and third molars perforate below the mylohyoid insertion directly into the submandibular and sublingual spaces, culminating in Ludwig's angina [1,2,4,6]. The defining clinical profile of this neglected state includes wooden induration of the submental region, severe trismus, marked dorsal elevation of the tongue, stridor, and precipitous asphyxiation [1,2,4,6]. When treatment for lower molar or pericoronal infections is withheld longer, purulent exudate traverses the lateral pharyngeal space into the danger space (Space 4), driving descending necrotizing mediastinitis with hallmark retrosternal chest pain, dyspnea, pleural empyema, and mortality rates ranging from 20% to 40% [1,2,6,7].

In susceptible hosts—particularly those with poorly controlled diabetes mellitus—unattended pulpal or periodontal necrosis easily triggers craniomaxillofacial necrotizing fasciitis along superficial and deep cervical fascial sheets, manifesting as rapid gangrenous necrosis, soft-tissue crepitus, and a lethal outcome in up to 30.3% of diabetic patients [5,6,7]. In the upper arch, periapical abscesses of the canines, premolars, and molars propagate through the valveless facial and angular veins and the pterygoid venous plexus, culminating in septic cavernous sinus thrombosis characterized by chemosis, proptosis, progressive ophthalmoplegia (cranial nerves III, IV, VI), and blindness [6,7]. Simultaneously, chronic occult apical or periodontal foci serve as insidious reservoirs for brain abscesses and subdural empyemas, where patients deteriorate with altered consciousness, focal deficits, and coma [6,7,8]. Chronic suppurative periodontitis and apical neglect disrupt osseous microcirculation, driving jaw osteomyelitis, extensive cortical bone sequestration, pathological mandibular fractures, and condylar necrosis leading to permanent temporomandibular joint ankylosis [1,6]. Finally, erosion of multi-space suppuration into the carotid sheath triggers septic internal jugular vein thrombophlebitis and bacteremia, resulting in septic pulmonary embolization, refractory septic shock, disseminated intravascular coagulation, and multiple organ failure [2,4,6,7].

REFERENCES

1. Pesis, M., Bar-Droma, E., Ilgiyaev, A., & Givol, N. (2019). Deep neck infections are life threatening infections of dental origin: A presentation and management of selected cases. *Isr Med Assoc J*, 21(12), 806–811.
2. Dalla Torre, D., Brunold, S., Kisielewsky, I., Kloss, F. R., & Burtscher, D. (2013). Life-threatening complications of deep neck space infections. *Wien Klin Wochenschr*, 125(21–22), 680–686.
3. Flynn, T. R., Shanti, R. M., Levi, M. H., Adamo, A. K., Kraut, R. A., & Trieger, N. (2006). Severe odontogenic infections, part 1: Prospective report. *J Oral Maxillofac Surg*, 64(7), 1093–1103.
4. Kataria, G., Saxena, A., Bhagat, S., Singh, B., Kaur, M., & Kaur, G. (2015). Deep neck space infections: A study of 76 cases. *Indian J Otolaryngol Head Neck Surg*, 67(Suppl 1), 83–88.
5. Gore, M. R. (2018). Odontogenic necrotizing fasciitis: A systematic review of the literature. *BMC Ear Nose Throat Disord*, 18, 14.
6. Bali, R. K., Sharma, P., Gaba, S., Kaur, A., & Ghanghas, P. (2015). A review of complications of odontogenic infections. *Natl J Maxillofac Surg*, 6(2), 136–143.
7. Evsen, E. A., & Candan, M. (2023). Serious complications and treatment strategies associated with odontogenic infections. *Eurasian J Med*, 55(1), 142–149.
8. Burgos-Larraín, L. F., Vázquez-Portela, Á., Cobo-Vázquez, C. M., Sáez-Alcaide, L. M., Sánchez-Labrador, L., & Meniz-García, C. (2022). Brain complications from odontogenic infections: A systematic review. *J Stomatol Oral Maxillofac Surg*, 123(6), e794–e800.